Eventlabel	JUB2000_sel_a_m_06
Campaign	King George Island 2000
Species	Southern elephant seal ( <i>Mirounga leonina</i> )
Age	≥6 years, adult
Sex	Male
Number	06
Length	390 cm
Girth	
Weight [estimated]	
Weight [calculated]	
Weight [measured]	
ARGOS PTT ID	16962
Transmitter type	SDR-T6, Half-Watt, Microprocessor-controlled Satellite-linked Time-Depth Recorder
Manufacturer	Wildlife Computers
PTT Serial Number	96-248
PTT Software	3.14a
Setting protocol	Half-Watt, Microprocessor-controlled Satellite-linked Time-Depth Recorder.  Unit measures depth from 0 to 1488 meters with a resolution of 6 meters  Software version 3.14a. Unit number: 96-248. ARGOS geolocation id = 16962  Unit identifier = JUB2000_sel_a_m_06. Unit started at 11:40:20 on 13/03/:0  Time (GMT) is 11:44:45.21. Date (GMT) is 13 March 19:0  Shallowest depth to be considered a "dive" = 12 meters  Deepest depth for accumulating surface-timelines (0=dry only) = 6 meters  SLTDR uses 1-sec / ¼-sec wakeups when shallower than 36 / 12 meters  Local time [0-23 hours] corresponding to 00h UT (GMT): 20  Transmission intervals (at-sea / on-land) = 00:51.00 / 01:32.00  SLTDR will use on-land interval after 10 consecutive dry transmissions  SLTDR will suspend transmissions after 6 hours "hauled-out". "Haul-out" ends after SLTDR is "wet" for 3 successive at-sea transmission intervals

	accumulate) = 300 STATUS will be transmitted every 24 messages. Blocks of Time-Lines will be transmitted every 48 messages. Hours when SLTDR transmits: 00-23,□ Upper limits of maximum-depth histogram bins are: 102, 204, 300, 402, 504, 600, 702, 804, 900, 1002, 1104, 1200, 1302, ∞ meters Upper limits of dive-duration histogram bins are: 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, ∞ minutes Upper limits of time-at-depth histogram bins are: 102, 204, 300, 402, 504, 600, 702, 804, 900, 1002, 1104, 1200, 1302, ∞ meters **** Check these parameters carefully ****. Ready to deploy? y Type D to archive depth readings, H to archive histograms: d
Deployment	
.,,,	Head, antenna cranial (45°)
Immobilisation	Large Animal Immobilon (LA Immobilon) was injected remotely by Telinject®-vario darts to achieve initial sedation (x=0.0009 mg/kg etorphine; 0.0037 mg/kg acepromacine; n= 27) while ketamine was injected manually on demand to maintain narcosis (x=81 min). The total dosages (x=1.7 mg/kg) of ketamine required were negatively correlated with those of LA Immobilon (p < 0.01). The dosages of LA Immobilon were approximately 15 to 30 times lower than recommended for other large-sized mammal species, and the therapeutic range was low. Nine cases required the application of the etorphine-antidote Large Animal Revivon (x=0.0052 mg/kg diprenorphine) injected intraveneously (n=3), intramuscularly (n=5), or sublingually (n=1).  Ramdohr, S., Bornemann, H., Plötz, J., Bester, M.N. (2001). Immobilisation of free-ranging adult male southern elephant seals ( <i>Mirounga leonina</i> ) with Immobilon (etorphine/acepromacine) and ketamine. South African Journal of Wildlife Research 3/4:135-140
Comment	3, 1123 210
Tag deployed	2000-03-18, -62.233, -58.667
Tag retrieved	
First transmission	2000-03-18T21:23:02, -62.312, -58.649
Last transmission	2000-10-26T10:22:05, -63.699, -56.901
	1